Draft version September 21, 2012
Preprint typeset using LATEX style emulateapj v. 12/16/11

arXiv:1209.4418v1 [astro-ph.EP] 20 Sep 2012

ON THE DIRECT IMAGING OF TIDALLY HEATED EXOMOONS

Mary Anne Peters[1,2]* and Edwin L. Turner[1,3]
[1] Department of Astrophysical Sciences, Princeton University, Princeton, NJ 08544, USA
[2] Department of Mechanical and Aerospace Engineering, Princeton University, Princeton, NJ 08544, USA and
[3] The Kavli Institute for the Physics and Mathematics of the Universe, The University of Tokyo, Kashiwa 227-8568, Japan

Draft version September 21, 2012

ABSTRACT

We demonstrate the ability of existing and planned future telescopes, on the ground and in space, to directly image tidally heated exomoons orbiting gas-giant exoplanets. Tidally heated exomoons can plausibly be far more luminous than their host exoplanet and as much as 0.1% as bright as the system's stellar primary if it is a low mass star. Because emission from exomoons can be powered by tidal forces, they can shine brightly at arbitrarily large separations from the system's stellar primary with temperatures of several hundreds degrees Kelvin or even higher in extreme cases. Furthermore, these high temperatures can occur in systems that are billions of years old. Tidally heated exomoons may thus be far easier targets for direct imaging studies than giant exoplanets which must be *both* young and at a large projected separation (typically at least tens of AU) from their primary to be accessible to current generation direct imaging studies. Specifically, current ground and space based instruments are capable of detecting exomoons with brightness temperature $\geq 600K$ and $R \geq 1R_\oplus$ in K-band. Future mid-infrared space telescopes, such as JWST and SPICA, will be capable of directly imaging tidally heated exomoons around the nearest two dozen stars with brightness temperature $\geq 300K$ and $R \geq 1R_\oplus$ orbiting at $\geq 12AU$ around stars within 4 parsecs of Earth at a 5σ confidence level in a 10^4 second integration. In addition it is possible that some of the exoplanets which have already been directly imaged are actually tidally heated exomoons or blends of such objects with hot young planets; we speculate that Fomalhaut b could be such a case. If such exomoons exist and are sufficiently common (*i.e.,* nearby), it may well be far easier to directly image an exomoon with surface conditions that allow the existence of liquid water than it will be to resolve an Earth-like planet in the classical Habitable Zone of its primary.

1. INTRODUCTION

Direct imaging of exoplanets, especially those in the "Habitable Zone" (HZ), is extremely difficult because of the high contrast ratio between the star and planet and because the of the very small star-planet angular separation. Indeed, all exoplanets that have been directly imaged to date are well separated from their host star, and are young systems that are still hot ($T_{eff} \sim 1000K$) from their formation (as opposed to being heated by stellar insolation). Examples include the HR8799 planets, β Pic b and LkCa15b (Marois et al. (2008), Lagrange et al. (2008) and Kraus & Ireland (2011)).

Although there has already been substantial discussion of the possibility of tidally heated exomoons (THEMs), extrasolar analogies of Solar System objects such as Io, Europa and Enceladus, in the literature (Peale et al. (1979), Yoder & Peale (1981), Ross & Schurbert (1987), Ross & Schubert (1989), Nimmo et al. (2007)) and even of their potential astrobiological interest ((Scharf (2006), Henning et al. (2009),Heller (2012)), the possibility of detecting exomoons has so far been restricted to indirect methods (Sartoretti & Schneider (1999), Han & Han (2002), Simon et al. (2007), Kipping (2009a), Kipping (2009b)).

This paper investigates the possibility that THEMs could be directly imaged (and perhaps already have been) with existing ground and space based instrumentation and even more effectively with currently planned future direct imaging facilities. *This scenario has several powerful potential advantages. THEMs may remain hot and luminous for periods of order a stellar main sequence lifetime and so could be visible around old stars as well as young ones. In addition, since THEMs may be hot even if they receive negligible stellar insolation, they may be luminous at large separations from the system primary, thus reducing or eliminating the requirement of high contrast imaging capabilities.* Moreover, tidal heating depends so strongly on of the orbital and physical parameters of an exomoon, that quite plausible systems (*i.e.,* with properties not very different from those occurring in the Solar System) will result in terrestrial planet sized objects with effective temperatures as high as $1000K$, or even higher in extreme but physically permissible cases.

In order to provide context and motivation for the analysis to follow, it is helpful to consider Solar System tidally heated moons. Io emits more energy per unit area at $\lambda \sim 5\mu m$ than expected (Witteborn et al. (1979), Spencer et al. (2005)) and has the highest measured temperatures of any body in the outer Solar System due to tidal heating (McEwen et al. (1997)). If the Galilean moon system orbited Neptune with the semi-major axes of their orbits scaled down in proportion to the Roche radius of that planet (relative to Jupiter's), Io would be more luminous than Neptune. If Io was as massive and dense as Earth, it would be the brightest Solar System object beyond 5 AU, out-shining even Jupiter in the 2-

---

*correspondence: mapeters@princeton.edu

$4\mu m$ and 5.5-6.5μm wavelength ranges[1]. Furthermore, if Io were as massive as Earth, it would be bright enough for JWST to detect it at a distance of 5 parsecs!

The remainder of this paper is organized as follows: We establish the exomoon tidal heating equations and present associated scaling relations in section 2. In section 3 we discuss discovery space constraints and determine the detection limits and observability of THEMs with existing and future instrumentation. Section 4 briefly presents a speculation that the SED of Fomalhaut b contains a significant contribution from a THEM. Conclusions and implications are described in section 5.

2. TIDAL HEATING

Tidal heating of moons in the Solar System, such as Io and Europa, have been analyzed in detail by Reynolds et al. (1987), Segatz et al. (1988) and S. J. Peale (1978). In this section we adapt the resulting equations for tidal heating of exomoons from these literature analyses. From these inputs the relevant scaling relations based on orbital, exomoon, exoplanet and host-star parameters are easily obtained.

2.1. *Luminosities*

Reynolds et al. (1987) and Segatz et al. (1988)) show that the average total luminosity of a moon due to tidal heating, L_{tidal} is given by

$$L_{tidal} = \frac{42\pi G^{5/2}}{19}\left(\frac{R_m^7 \rho^2 M_p^{5/2}}{\mu Q}\right)\left(\frac{e^2}{a^{15/2}}\right) \tag{1}$$

where G is the gravitational constant, μ is the moon's elastic rigidity, Q is the moon's dissipation function (or quality factor), e is the eccentricity of the moon's orbit, a is the semi-major axis of the moon's orbit, ρ is the density of the moon, R_m is the radius of the moon, and M_p is the mass of the planet it orbits.

It is useful to eliminate the explicit dependence on the planet's mass by parameterizing equation 1 in terms of the Roche radius. Let the moon's semi-major axis be some multiple, β, of the Roche radius, a_R, such that

$$a = \beta a_R = \beta\left(\frac{3M_p}{2\pi\rho}\right)^{1/3}. \tag{2}$$

Then we can rewrite the tidal energy flux equation as

$$L_{tidal} = \left(\frac{6272\pi^7 G^5}{9747}\right)^{1/2}\left(\frac{R_m^7 \rho^{9/2}}{\mu Q}\right)\left(\frac{e^2}{\beta^{15/2}}\right) \tag{3}$$

where we have grouped the terms that depend on the moon's physical properties and those that describe its orbit separately.

Alternatively, we can write this equation in terms of R_m and M_m or ρ and M_m rather than R_m and ρ. Alternative forms of the Eq. 3 are

$$L_{tidal} = \left(\frac{189 G^{5/2}}{608\sqrt{2}\pi}\right)\left(\frac{M_m^{9/2}}{\mu Q R_m^{13/2}}\right)\left(\frac{e^2}{\beta^{15/2}}\right) \tag{4}$$

$$L_{tidal} = \frac{7}{19}\left(\frac{243\pi^7 G^{15}}{128}\right)^{1/6}\left(\frac{M_m^{7/3}\rho^{13/6}}{\mu Q}\right)\left(\frac{e^2}{\beta^{15/2}}\right) \tag{5}$$

The scaling relation for equation 3 relative to the luminosity of Earth (which is $L_\oplus \sim 1.75\times10^{24}$ ergs/s) is

$$L_m \approx L_\oplus\left[\left(\frac{R_m}{R_\oplus}\right)^{\frac{5}{4}}\left(\frac{\rho}{\rho_\oplus}\right)^{\frac{9}{8}}\left(\frac{36}{Q}\cdot\frac{10^{11}\frac{dynes}{cm^2}}{\mu}\right)^{1/4}\right] \times\left[\left(\frac{e}{0.0035}\right)^{\frac{1}{2}}\left(\frac{\beta}{8}\right)^{-\frac{15}{8}}\right]. \tag{6}$$

Note that the first set of bracket terms in Eq. 3 includes exomoon's physical parameters, and the second contains the orbital parameters. This scheme is also used in equations 8 and 9 below.

2.2. *Effectives Temperatures and SEDs*

Using the normal conventions of stellar astrophysics, we can define the exomoon's effective temperature, T_m, from the luminosity via the Stefan-Bolzmann Law

$$T_m = \left(\left(\frac{392\pi^5 G^5}{9747\sigma^2}\right)^{1/2}\left(\frac{R_m^5 \rho^{9/2}}{\mu Q}\right)\left(\frac{e^2}{\beta^{15/2}}\right)\right)^{1/4}. \tag{7}$$

Eq. 7 can also be written as a scaling relation relative to a 300K exomoon.

$$T_m \approx 300K\left[\left(\frac{R_m}{R_\oplus}\right)^{\frac{5}{4}}\left(\frac{\rho}{\rho_\oplus}\right)^{\frac{9}{8}}\left(\frac{36}{Q}\cdot\frac{10^{11}\frac{dynes}{cm^2}}{\mu}\right)^{1/4}\right] \times\left[\left(\frac{e}{0.004}\right)^{\frac{1}{2}}\left(\frac{\beta}{8}\right)^{-\frac{15}{8}}\right] \tag{8}$$

Note that equation 8 adopts $Q = 36$ and $\mu = 10^{11}$ dynes/cm^2 as reference values. These are the Q and μ values for Io from the literature (Segatz et al. (1988) and Peale et al. (1979)). Equation 8 also assumes there is only tidal heating with no additional energy sources, such as stellar insolation or interior radiogenic heat. This is a conservative assumption since additional heat sources only serve to make the exomoon more luminous and thus easier to detect. The scaling relations in Eq. 8 are illustrated in Fig. 1.

Given T_m, the peak wavelength of a exomoon's spectral energy distribution (SED) would be

$$\lambda_{max} \approx 9.6\mu m\left[\left(\frac{R_m}{R_\oplus}\right)^{-\frac{5}{4}}\left(\frac{\rho}{\rho_\oplus}\right)^{-\frac{9}{8}}\left(\frac{36}{Q}\cdot\frac{10^{11}\frac{dynes}{cm^2}}{\mu}\right)^{1/4}\right] \times\left[\left(\frac{e}{0.004}\right)^{-\frac{1}{2}}\left(\frac{\beta}{8}\right)^{\frac{15}{8}}\right] \tag{9}$$

if it emitted as an ideal blackbody. However, Solar System objects with significant tidal heating typically do not have a uniform temperature surface emitting at

---

[1] Based on Spiegel & Burrows (2012) model of a 1M$_J$ 1Gyr old cloud-free, solar metalicity Jupiter-like planet, and assumes a blackbody curve for the scaled up version of Io.



FIG. 1.— LEFT: Plot of eccentricity vs. effective temperature of an exomoon, with a semi-major axis of 5 Roche radii (for comparison, Io is at 6.6 Roche radii). The solid and dotted lines correspond to a density of $\rho = 3.5 g/cm^3$ and $\rho = 5.5 g/cm^3$ respectively. The lower density ($\rho = 3.5 g/cm^3$) matches that of Io. The higher density ($\rho = 5.5 g/cm^3$) matches Earth. Each line color corresponds to a different moon radius listed in the upper right corner of the plot on the right (note that this legend lists the moon radii for both plots). RIGHT: Plot of semi-major orbital axis vs. effective temperature of an exomoon. This plot assumes a moon with eccentricity of 0.005 (for comparison, Io has an eccentricity of $e = 0.004$). Dotted and solid lines represent the same densities as the plots on the left.

each point as an ideal blackbody. Rather they display "hot spots" and even vulcanism, locations on the surface through which a larger, often much larger, than average part of the internal heating is being radiated away. This implies an SED that deviates significantly from a Planck form and which, in particular, emits more at shorter wavelengths than the uniform temperature blackbody approximation indicates. This is seen in the SED of Io, for example (Spencer et al. (2005)). However, modeling of the complexities of heat transport in the interior of a THEM far exceeds the scope of this initial discussion of detectability, and thus we hereinafter adopt the simple blackbody model of THEM SEDs. In most respects, this is a conservative assumption in that it makes them less detectable than they would be expected to be with a more realistic SED. Note that the blackbody SED assumption will be more accurate for an exomoon on which a thick atmosphere and/or oceans effectively redistribute the tidal heat emitted at hot spots on its surface.

2.3. *Contrast*

Discussions of direct imaging of exoplanets are typically heavily focused on issues of contrast and associated instrumental inner working angles for the star-exoplanet separation on the plane of the sky. The associated considerations for THEMs are more complex since both contrast with the planet orbited by the exomoon and contrast with the stellar primary must be taken into account. Since the star-exoplanet separation does not affect the tidal luminosity of a given exomoon system, the contrast with the star may not (or may) be an important observational issues. However, since tidal heating is very sensitive to the moon-planet separation, it is not plausible that a significantly bright THEM can be resolved from the planet it orbits with existing or planned facilities. In other words, any emission from an exoplanet will dilute that from any THEM which orbits it.

Tidally heated moons are easier to detect if L_m is large and L_p and L_* are small. Equation 3 gives the tidal luminosity of an exomoon. The contrast of the moon with the planet (L_m/L_p) decreases for colder exoplanets ($L_p \propto T_p^4$) that are further away from their host star and for older exoplanets which have already cooled from their initial formation temperatures. The moon-planet contrast also decreases for an exoplanet with less surface area ($L_p \propto R_p^2$), however, larger exoplanets provide more gravitational potential energy and are thus able maintain a THEM's temperature for a longer period of time. The contrast of the moon relative to the star (L_m/L_*) decreases for less massive stars ($L_* \propto M_p^{2.3}$ for $M < 0.43 M_\odot$; Burrows et al. (2001)). Finally, the contrast requirement will be relaxed for relatively nearby star systems due to the resulting larger angular separations on the sky.

2.4. *Lifetime*

TABLE 1
Relevant physical parameters for various solar system bodies

Parameter[a]	Io	Europa	Enceladus	Moon	Titan	Earth
a (10^3km)	422	671	238	384	1222	-
β	6.63	10.0	3.82	40.5	23.3	-
e	0.0041	0.0101	0.0045	0.055	0.029	-
R (km)	1821	1560	250	1737	2575	6371
ρ (g/cm^3)	3.53	3.02	1.61	3.35	1.88	5.52
μ (10^9 $\frac{dynes}{cm^2}$)	(100)[b]	(40)	(2000)	50	(9)	1200
Q	(36)	(100)	(100)	27	(100)	280

[a] References for values are Lodders & Fegley (1998), Segatz et al. (1988), Scharf (2006), Murray & Dermott (1999), Yoder (1995), Thomson (1863) and Ray et al. (2001)

[b] Parenthesis indicate this is a modeled, not measured value.

In order to sustain tidal heating, a moon's orbital eccentricity must be maintained by resonance with another moon. For example, Io, Europa and Ganymede are in a 1:2:4 orbital period resonance. This resonance preserves Io's orbital eccentricity and has sustained Io's tidal heating over a time of comparable to the lifetime of the Solar System. Without the resonance the orbit and eccentricity of a tidally heated moon will decay in the same fashion as the lunar eccentricity (and angular momentum) have decayed over time. Hence, we are only likely to find tidally heated exomoons in systems where they are in resonance with other exomoons, however there is no reason to believe that such resonances are uncommon. Both Jupiter and Saturn have close-in moons that are in resonance (Peale (1976)), and there are theoretical reasons to expect such resonances to develop naturally during the formation process (Yoder (1979), Cassidy et al. (2009), Ogihara & Ida (2012)).

In general, more massive planets and more massive moons are expected to give lead to longer lifetimes for the orbital resonances that sustain the tidal heating, at fixed semi-major axis measured in Roche radii.

2.5. *Variability*

In general the observed brightness and SED of a THEM is expected to be substantially variable for multiple reasons: Most dramatically the exomoon may be eclipsed by the much larger and darker exoplanet it orbits, thus causing a sharp drop (and later increase) in the observed flux. The steep dependence of tidal heating on semi-major axis implies that the most luminous exomoons will have close in orbits and thus particularly large eclipse probabilities. Even in the absence of eclipses, phase curve variations are expected since tidal heating typically produces moderate to extreme temperature variation across an object's surface. Moreover, even at a single location on the exomoon's tidally heated surface, the temperature may well fluctuate due to time varying transport of interior heat to the surface, *e.g.,* vulcanism. In general short timescale (hours to days would be expected from Solar System analogs) variability would be a signature of THEMs which would help distinguish them from the relatively steady emission expected from a cooling gas-giant exoplanet.

3. DETECTABILITY

3.1. *Temperature Limit*

Before exploring the THEM discovery space it is necessary to understand constraints on exomoon temperatures that can be produced by tidal heating. The temperature power law dependancies are so strong that the simple scaling relations presented in the previous section yield temperatures of thousands of degrees for seemingly plausible hypothetical exomoon systems. However, there are other physical constraints that limit the effective temperatures that can be achieved by tidal heating. These constraints are discussed in Cassidy et al. (2009). We note that long-lived resonances in the solar system exist, and thus should exist in other systems as well.

The Cassidy et al. (2009) analysis does not yield any precise upper limit on the surface temperature of an exomoon; we therefore conservatively adopt $T_m = 1000K$ in our analysis. As context for this number note that the surface temperatures of known rocky bodies, such as Mercury and UCF-1.01 in the GJ 436 system, are of this order. The Sun facing side of Mercury reaches temperature of 700K, and UCF-1.01's surface is estimated to be at $\sim 860K$ (Stevenson et al. (2012)). If rocky planets can survive surface temperatures near 1000K, rocky exomoons should be able to do so as well.

3.2. *Assumed Exomoon Properties*

The quantities Q and μ are poorly known even for most Solar System objects. For the first-approximation models considered here, we adopt the values for Io in the literature. Table 1 lists relevant quantities for several moons in the Solar System as well as the Earth. Io is a rocky moon with a hot, molten core which is probably the most relevant comparison for a strongly heated exomoon. For parameters other than Q and μ, we assume that the range of values shown in Table 1 indicates plausible cases.

3.3. *Existing Facilities*

Fig. 2 illustrates the detectability of tidally heated exomoons with existing ground and space-based facilities. The exomoon curves are shown as blackbodies with temperatures defined by Eq. 8 and with the constraints and assumptions discussed in the previous sections. The curves shown for the three stars are from the Kurucz models (Kurucz (1970)). The red dotted line is the five times the *square-root* of the atmospheric emission at Mauna Kea, not the incident flux (*i.e.,* this is five times the $\sqrt{N}$ noise from the atmosphere, which roughly corresponds to the 5σ detection limit). This noise floor assumes the collecting area of a 8.1m telescope, a $100 \times 100 mas^2$ full-width half-max point spread function, a 10,000s integration time, and uses the Lord (1992) atmospheric transmission tool. Note



FIG. 2.— The top three lines are the energy flux received from HR8799 (black line), the sun (dark blue) and an M5v star (light blue) at a distance of 10 parsecs. The lines labeled with a temperature and radius are four plausible exomoon blackbody curves. Note the Io is approximately one quarter the radius of Earth. The light gray line is modeled exoplanet 1M$_J$ in mass, 1Gyr old and with no clouds and solar metalicity (Spiegel & Burrows (2012)). Note that this is younger (*e.g.* brighter) than most stars in the solar neighborhood. The dotted red line is the 5σ noise floor due to atmospheric emission. Left-most contrast line shows the contrast of HR8799b relative to it's host star in the K-band. The middle contrast line shows the contrast of a 1000K, one Earth mass moon relative to a sunlike-star in the K-band. The right contrast line shows the same, but in L-band where the star-exomoon contrast is less.

that this curve represents an idealized case with 100% transmission optics. Current ground based instruments will be limited by atmospheric emission beyond the K-band. The high contrast and adaptive optic (AO) systems currently on the infrared platform at the Subaru telescope, namely HiCIAO (Hodapp et al. (2008)) and AO188 (Hayano et al. (2010), Minowa et al. (2010)), are currently capable of achieving 10^{-5} contrast at 0.2 arcsec angular separation from the host star. The implementation of the next generation of high contrast instrumentation, SCExAO (Martinache et al. (2011)) and CHARIS (McElwain et al. (2012)) will be able to do an order of magnitude better than that (10^{-6} contrast) at the same separation, and 10^{-7} contrast at a 2 arcsec separation. Existing space based instruments on HST and Warm Spitzer such as Spitzer's IRAC have similar detection limits to ground based observatories (see Spitzer detection limits in Fig. 2). However, HST and Spritzer do not have the atmospheric transmission and turbulence issues associated with ground based observing, which may make spaced-based detection of exomoons easier in comparison with existing ground based facilities.

The easiest detection of an exomoon with ground-based instrumentation would occur in the L-band. Fig. 2 shows that a 1000K, one Earth-radius exomoon would be less than 10^3 times dimmer than a late-type M-dwarf at these wavelengths. Telescopes equipped with adaptive secondaries and adaptive optics operating in the L-band may be optimal for detecting these moons from the ground. The high contrast planet imagers (such as GPI, SPHERE and CHARIS) (Macintosh et al. (2008), Beuzit et al. (2008), Peters et al. (2012)) coming on sky in the next 1-3 years should be able to detect 600K, one Earth radius moons in the K-band in systems with late-type M-dwarfs at a distance 5 parsecs. This same exomoon would have even more favorable contrast with it's host star at $\lambda = 4.5\mu m$ with warm Spitzer, but this detection



Fig. 3.— The top two blue lines are the energy flux received from the sun (dark blue, top) and M5v star (light blue) at a distance of 3 parsecs. The lines labeled with a temperature and radius are four plausible exomoon blackbody curves. The gray line is the 1Gyr, $1M_J$ planet in Fig. 2, but scaled to the luminosity of Jupiter so that it represents an older planet, more typical of the ages of planets found in the local neighborhood. Left contrast line shows that a 300K, Earth-radius moon would be $10^4\times$ fainter than the M5v star. The contrast line on the right shows that a 300K, $R = 2R_\oplus$ object would be $10^5\times$ fainter than a sun-like and $\sim 5\times$ brighter than a Jupiter. Note that the x- and y-axis are different than in Fig. 2.

would only be possible if the exomoon was at a distance of 1 parsec. It is interesting to note that at these temperatures, the exomoon would likely be many orders of magnitude brighter than its host planet at most wavelengths, even if the planet were 10× more massive than Jupiter, assuming a system with an age comparable to the Solar System's.

3.4. *Future Facilities*

Future instruments, for example JWST's Mid-Infrared Instrument (MIRI), offer even more promising potential for direct imaging of exomoons. Fig. 3 shows the discovery space for MIRI. Similar to Fig. 2, the solid lines labeled with temperatures and radii correspond to exomoon blackbody curves with those parameters. Note that the exomoons shown here have temperatures similar to Earth's rather than the much hotter temperatures shown in Fig. 2. Thus, it is plausible that some of the exomoons JWST is capable of detecting, could potentially be habitable, in the sense of having surface temperatures that would allow liquid water to be present. Some of these exomoons have comparable irradiance to the gas giants in our solar system. At $\sim 14\mu m$ a 300K, an Earth-radius exomoon would be as luminous as Jupiter. However, if Jupiter were colder due to being less heated by its primary and/or being older, the Earth-like moon would be much brighter than the planet. The red bars are the 5σ detection limits for JWST-MIRI with a 10,000s integration time (Glasse et al. (2010)).

A 300K, Earth-radius tidally-heated exomoon is only 3×10^5 times fainter at $\lambda \sim 14\mu m$ than a Sun-like star, but can be at a large distance from it's host star. At 30AU projected separation, it would be 15 arcsecs from the star if it were 2 parsecs away. At $\lambda = 14\mu m$, λ/D=0.44 arcsecs for a 6.5m telescope, which means this moon would be at $30\lambda/D$. This far away from the star, the airy rings are $\sim 3 \times 10^5$ times fainter than the core of the star and about the same intensity as the 300K, Earth-radius moon, indicating that the detection should be possible. If there is such an Earth-sized 300K moon orbiting αCen, MIRI will be able to detect it in 8 of its 9 spectral bands with better than 15 sigma signal-to-noise. Thus, directly imaging a 300K, Earth-radius moon that is tidally heated is potentially much easier than resolving an Earth-like exoplanet orbiting in the HZ of its primary.

SPICA is another future space telescope that is ideal for exomoon detection. The SPICA coronagraph is being designed to operate from $3.5 - 27\mu m$ with a contrast of order 10^{-6} and a $3.3\lambda/D$ inner working angle (Enya et al. (2011)). SPICA has a slightly higher noise floor than JWST, but should be able to achieve similar contrast. Thus SPICA's exomoon discovery space will be similar to JWST's, though JWST's larger aperture will give MIRI a modest advantage.

The next generation of ground based telescopes such as the Giant Magellan Telescope (GMT, Johns (2008)), the Thirty Meter Telescope (TMT, Nelson & Sanders (2008)), and the European Extremely Large telescope (E-ELT, McPherson et al. (2012)) are best suited for detecting THEMs at closer exomoon-star separations and at slightly shorter wavelengths than SPICA and JWST. The sensitivity limits of the Mid-Infrared E-ELT Imager and Spectrograph (METIS) will be almost 10× better than Spitzer's detection limit (Brandl et al. (2010)). The high contrast exoplanet imagers, such as E-ELT's EPICS (Kasper et al. (2010)), claim they will be able to achieve contrasts of 2×10^{-10} at 0.2 arcsec angular separations.

4. FOMALHAUT B

Fomalhaut is an A4v young (200-400 Myrs) main sequence star at a distance of 7.7 pc orbited by a bright and extensive circumstellar disk. In 2008, Kalas et al. (2008) reported an exoplanet detection around Fomalhaut in the F606W and F814W HST bands (~500-1000nm range). However, subsequent infrared observations between 1.5-$8\mu m$ and non-detections suggested that this object is not an exoplanet. In addition, the emission in the F606W band varied substantially between observations taken in 2004 and 2006 (Miller-Ricci et al. (2009), Marengo et al. (2009), Janson et al. (2012)). Alternative explanations have been given for the object including the possibility of transient dust clouds (Janson et al. (2012)) and planetesimal bombardment heating of the exoplanet's surface (Miller-Ricci et al. (2009)).

We propose a new possibility, namely that Fomalhaut b is a THEM or, perhaps more plausibly, a blend of emission from a young and hot gas-giant exoplanet and a THEM. A sufficiently hot and small THEM could explain the lack of emission in the infrared and the excess emission in the F606W band as well as helping explain the apparent brightness variations. Eclipses of this hypothetical exomoon would occur if the exomoon's orbit has $a < 2.3R_{planet}$ assuming its orbit is coplanar with the circumstellar disk.

Given the limited available photometry and the many available free parameters, it seems certain that a blend model, combining the light of a THEM with that of a hot young gas-giant could provide an acceptable fit to the data. However, such modeling is beyond the scope of the present paper. More promisingly, additional observations might well be able to confirm or strongly support the THEM hypothesis for Fomalhaut b. Further observations of this system with HST could be conclusive in themselves if the planet is found to eclipse the exomoon, causing the flux at short wavelengths to disappear regularly on a plausible orbital period.

5. CONCLUSIONS

Direct imaging detection of physically plausible, tidally heated exomoons is possible with existing telescopes and instrumentation. If tidally heated exomoons are common, for example if typical gas giant exoplanets are orbited by satellite systems broadly similar to those found in the Solar System, we are likely to be able to image them around nearby Sun-like stars in the midst of their main sequence lifetimes with current or near future facilities.

Existing instrumentation should be able to detect exomoons with temperatures $\geq 600K$ and $R \geq 1R_\oplus$ in K-band at 5 parsecs. Future mid-infrared space telescopes such as JWST and SPICA will be capable of directly imaging tidally heated exomoons around the nearest two dozen star systems with brightness temperature $\geq 300K$ and $R \geq 1R_\oplus$ orbiting at $\geq 12AU$ around stars within 4 parsecs of Earth at the 5σ confidence level in multiple bands. It is possible that some of the exoplanets which have already been directly imaged, such as Fomalhaut b, could be tidally heated exomoons or exoplanet-exomoon blends. It is therefore plausible that a habitable (in the sense of possessing liquid water on its surface) exomoon can be imaged long before it will be possible to do so for a habitable (in the same sense) exoplanet.

Thus, the era of astrobiology based on direct imaging of extrasolar objects may not have to await the advent of specialized space-based telescopes such as those contemplated for the TPF and DARWIN missions.

ACKNOWLEDGEMENTS

We thank Alexis Carlotti, Jill Knapp, Matt Mountain, George Rieke, Dave Spiegel and Scott Tremaine for useful conversations and comments. This research has been supported in part by World Premier International Research Center Initiative, MEXT, Japan.

REFERENCES

Beuzit, J., Feldt, M., Dohlen, K., et al. 2008, in Proc SPIE, Vol. 7014, 41

Brandl, B., Blommaert, J., Glasse, A., Lenzen, R., & Pantin, E. 2010, Msngr, 140, 30

Burrows, A., Hubbard, W., Lunine, J., & Liebert, J. 2001, Rev. Mod. Phys., 73, 719

Cassidy, T. A., Mendez, R., Arras, P., Johnson, R. E., & Skrutskie, M. F. 2009, ApJ, 704, 1341

Enya, K., Kotani, T., Haze, K., et al. 2011, AdSR, 48, 323

Glasse, A., Bauwens, E., Bouwman, J., et al. 2010, Proc SPIE, 7731, 77310K

Han, C., & Han, W. 2002, Astrophys.J., 580, 490

Hayano, Y., Takami, H., Oya, S., et al. 2010, Proc SPIE, 7736

Heller, R. 2012, In prep

Henning, W. G., O'Connell, R. J., & Sasselov, D. D. 2009, ApJ, 707, 1000

Hodapp, K. W., Suzuki, R., Tamura, M., et al. 2008, Proc SPIE, 7014, 701419

Janson, M., Carson, J., Lafreniere, D., et al. 2012, ApJ, 747, 116

Johns, M. 2008, Proc SPIE, 6986, 698603

Kalas, P., Graham, J. R., Chiang, E., et al. 2008, Science, 322, 1345

Kasper, M., Beuzit, J.-L., Verinaud, C., et al. 2010, Proc SPIE, 7735, 77352E

Kipping, D. M. 2009a, Mon. Not. R. Astron. Soc., 392, 181

—. 2009b, Mon. Not. R. Astron. Soc., 396, 1797

Kraus, A. L., & Ireland, M. J. 2011, ApJ, 745, 5

Kurucz, R. L. 1970, SAO Special Report no. 309

Lagrange, A. M., Gratadour, D., Chauvin, G., et al. 2008, A&A, 493, 5

Lodders, K., & Fegley, B. 1998, The Planetary Scientist's Companion (Oxford University Press)

Lord, S. D. 1992, A New Software Tool for Computing Earth's Atmospheric Transmission of Near- and Far-Infrared Radiation, Tech. rep., NASA Technical Memorandum 103957

Macintosh, B., Graham, J., Palmer, D., et al. 2008, in Proc SPIE, Vol. 7015, 7015–43

Marengo, M., Stapelfeldt, K., Werner, M. W., et al. 2009, ApJ, 700, 1647

Marois, C., Macintosh, B., Barman, T., et al. 2008, Science, 322, 1348

Martinache, F., Guyon, O., Garrel, V., et al. 2011, in Proceedings of SPIE, Vol. 8151, 81510Q

McElwain, M. W., Brandt, T. D., Janson, M., et al. 2012, Scientific Design of a High Contrast Integral Field Spectrograph for the Subaru Telescope

McEwen, A. S., Simonelli, D. P., Senske, D. R., et al. 1997, Geo. R. L., 24, 2443

McPherson, A., Gilmozzi, R., Spyromilio, J., Kissler-Patig, M., & Ramsay, S. 2012, The Msngr, 148

Miller-Ricci, E., Meyer, M. R., Seager, S., & Elkins-Tanton, L. 2009, ApJ, 704

Minowa, Y., Hayano, Y., Oya, S., et al. 2010, in Proc SPIE, Vol. 7736, 122

Murray, C. D., & Dermott, S. F. 1999, Solar System Dynamics (University of Cambridge)

Nelson, J., & Sanders, G. H. 2008, Proc SPIE, 7012, 70121A

Nimmo, F., Spencer, J. R., Pappalardo, R. T., & Mullen, M. E. 2007, Nature, 447, 289

Ogihara, M., & Ida, S. 2012, ApJ, 753, 60

Peale, S. J. 1976, ARAA, 13, 215

Peale, S. J., Cassen, P., & Reynolds, R. 1979, Science, 203, 892

Peters, M. A., Groff, T., Kasdin, N. J., et al. 2012, Conceptual Design of the Coronagraphic High Angular Resolution Imaging Spectrograph (CHARIS) for the Subaru Telescope

Ray, R. D., Eanes, R. J., & Lemoine, F. G. 2001, Geophys. J. Int., 144, 471

Reynolds, R. T., McKay, C. P., & Kasting, J. F. 1987, Adv. Space Res., 7, 125

Ross, M. N., & Schubert, G. 1989, Icarus, 78, 90

Ross, M. N., & Schurbert, G. 1987, Nature, 325, 133

S. J. Peale, P. C. 1978, Icarus, 36, 245

Sartoretti, P., & Schneider, J. 1999, A&A, 134, 553

Scharf, C. A. 2006, Astrophys.J., 648, 1196

Segatz, M., Spohn, T., Ross, M. N., & Schubert, G. 1988, Icarus, 75, 187

Simon, A., Szatmary, K., & Szabo, G. 2007, A&A, 470, 727

Spencer, J. R., Lellouch, E., Richter, M. J., et al. 2005, Icarus, 176, 283

Spiegel, D. S., & Burrows, A. 2012, ApJ, 745, 174

Stevenson, K. B., Harrington, J., Lust, N. B., et al. 2012, ApJ, 755, 9

Thomson, W. 1863, Phil. Trans. R. Soc. London, 153, 583

Witteborn, F. C., Bregman, J. D., & Pollack, J. B. 1979, Science, 203, 643

Yoder, C. F. 1979, Nature, 279, 767

—. 1995, Astrometric and geodetic properties of Earth and the Solar System (AGU Books Board)

Yoder, C. F., & Peale, S. J. 1981, Icarus, 47